

Severe Bronchial asthma induced by Gastro esophageal reflux disease – a case report

NS Neki^{1,*}, Satpal Aloona², Bhupinder Singh³

¹Professor, ^{2,3}Assistant Professor, Dept. of Medicine, Govt. Medical College & Guru Nanak Dev Hospital, Amritsar, Punjab

***Corresponding Author:**

Email: drneki123@gmail.com

Abstract

Gastroesophageal events remain an important factor in the pathogenesis of patients with asthma with the reported prevalence ranging from 32-82%. The esophageal bronchial reflux plays a greater role in the pathophysiology of reflex associated cough. Gastroesophageal reflux either singly or in association with postnasal drip and or asthma is a thought to be a cause of chronic cough. People with asthma are twice as likely to have GERD as those people who don't have asthma. GERD may worsen asthma symptoms, however asthma and some asthma medications may worsen GERD symptoms. Although these two disorders often occur together, the relationship between GERD and asthma remains unclear.

Keywords: Bronchial asthma; Gastroesophageal reflux disease(GERD); Proton pump inhibitors (PPIs).

Introduction

Gastroesophageal reflux disease (GERD) has been considered to be associated with cough in one third of patients.⁽¹⁾ Various studies have supported this documentation and shown a relationship between these two disorders.^(2,3) Chronic cough significantly affects the quality of life of these patients.⁽⁴⁾

Case Report

A 65 years old male presented with chronic cough, sputum and respiratory difficulty with repeated attacks of respiratory distress since 18 months having strong relationship with sleep and heavy meals. He was non-smoker, non-alcoholic with no H/O allergy. He had no family H/O bronchial asthma or allergy. He used to experience severe attacks of nocturnal wheezing, respiratory difficulty, orthopnea & tachypnea. These attacks used to subside with change of sleep posture and worsened with spicy food. The laboratory investigations in the form of CT scan, MRI, X-ray chest, cardiac evaluation & tracheoscopy did not reveal any abnormality. He was got examined by ENT specialist with diagnosis of allergic rhinitis without relief. With the passage of time, his symptoms started worsening & was admitted two times in the emergency department where he was managed with oxygen, bronchodilators & steroids. He was diagnosed to be a case of bronchial asthma but he started complaining of dyspepsia & fullness in the epigastium. He was subjected to be upper intestinal endoscopy which was normal. But esophageal manometry revealed lower LES relaxation of 6.5 mmHg (N 10-45 mmHg), low relaxation pressure of -20 mmHg (N <8 mmHg) with relax rate of 32% (N >80%). The patient was put on omeprazole and domperidone along with asthma medication. The intensity of symptoms subsided with no episodes of nocturnal awakening, cough &

asthmatic attacks. He was able to sleep with marked improvement in the quality of life.

Discussion

Extra esophageal presentations of GERD such as asthma, chronic cough, postnasal discharge & laryngeal disorders have been widely reported.⁽⁵⁾ In USA, 7-15% of western population has GERD, which is the 3rd most common GI disorder.⁽⁶⁾ There is limited data from Asia including India. 24 hour ambulatory esophageal pH monitoring and esophageal manometry are diagnostic of GERD without evidence of esophagitis. Takenaka R in their study⁽⁷⁾ using FSSG scale questionnaire revealed +ve prevalence of GERD in bronchial asthma as 37.4%. Both the esophagus and airways/lungs are vagally innervated & vice versa. This esophageal bronchial reflex is mainly responsible for gastroesophageal reflux leading to cough.⁽⁸⁾ The shared autonomic innervations is a consequence of the common origin of the esophagus and bronchial tree from the foregut.⁽⁹⁾ GERD is a potential trigger of asthma, although not all asthma patients with GERD experience reflex symptoms. Asoom AL, et al⁽¹⁰⁾ found that 36.4% of asthmatic patients diagnosed by esophageal pH monitoring as having GERD while Boma G et al⁽¹¹⁾ reported that about 40% of bronchial asthma patients had GERD. Legget J, et al in their study⁽¹²⁾ revealed that GERD is the main culprit for making asthma difficult to control. The association between GERD and pulmonary disease is further documented by the reduction or even disappearance of asthmatic symptoms after medical or surgical treatment of acid reflux⁽¹³⁾ as documented in our case. The treatment of GERD in asthma includes lifestyle modification & use of proton pump inhibitors (PPI) & surgical treatment.^(14,15)

Conclusion

Physicians should note that all severe asthma patients should be thoroughly investigated to exclude atypical presentation of GERD as in our case. GERD is a potential trigger of asthma although not all asthma patients with GERD experience reflux symptoms. The patients with asthma need to be put on PPIs and referred to gastroenterologist if need be.

References

1. Chung KF, Pavord ID. Prevalence, pathogenesis and causes of chronic cough. *Lancet* 2008;371:1364-74.
2. Smith JA, Decalmer S, Kelsall A, et al. Acoustic – Cough- Reflex Associations in chronic cough: Potential Triggers and Mechanisms. *Gastroenterology* 2010;139:754–762.
3. Kunch S, Gross V, Neesse A, et al. Combined lung sound and reflux-monitoring: a pilot study of a novel approach to detect nocturnal respiratory symptoms in gastroesophageal reflux disease. *Aliment Pharmacol Ther* 2011;33:592-600.
4. Young EC, Smith JA. Quality of life in patients with chronic cough. *Therapeutic advances in respiratory disease* 2010;4:49-55.
5. Poelmans J and Tack J. Extra-esophageal manifestations of gastroesophageal reflux. *Gut* 2005;54:1492-1499.
6. Triadafilopoulos G. GERD: The potential for endoscopic intervention. *Dig Dis* 2004;24:181-188.
7. Takenaka R. The use of FSSG scale in assessment of GERD in asthma. *Allergol Immunopathol* 2010;38(1):20-24.
8. Ing AJ, NguMc and Breslin AB. Pathogenesis of chronic persistent cough associated with gastroesophageal reflux. *Am J Respir Crit Care Med* 1994;149:160-167.
9. Richter JE. Gastroesophageal reflux disease and asthma: the two are directly related. *American Journal of Medicine* 2000;108(Suppl 1):153-158.
10. Asoom AL, Rubaish AM and Munshid HA. Gastroesophageal reflux in bronchial asthma patients. A clinical note. *Saudi Medical Journal* 2003;24(7):1364-1369.
11. Boma G and Gaude GS. Prevalence of gastroesophageal reflux disease in bronchial asthma patients. *Lung India* 2012;29:S22.
12. Leggett JJ, Johnston BT, Milles M, et al. Prevalence of gastroesophageal reflux in difficult asthma. Relationship to asthma outcome. *Chest* 2005;127(4):1227-1231.
13. Larrain A, Carroscio E, Galleguillos F, et al. Medical and surgical treatment to non-allergic asthma associated with gastroesophageal reflux. *Chest* 1991;99:1330-1336.
14. Bowrey DJ, Peters JH and Demeester TR. Gastroesophageal reflux disease in asthma: effects of medical and surgical anti-reflux therapy on asthma control. *Ann Surg* 2000;231(2):161-172.
15. Sharma B, Sharma M, Daga MK, et al. Effect of omeprazole and domperidone on adult asthmatics with gastroesophageal reflux. *World J Gastroenterol* 2007;13:1706-1710.